

AGGRESSION AND SELF-INJURY: RESEARCH NEEDS FOR THE SEVERELY AFFECTED END OF THE SPECTRUM

Matthew Siegel, M.D.

Associate Professor of Psychiatry & Pediatrics
Tufts University School of Medicine

VPMA, Developmental Disorders Service
Maine Behavioral Healthcare

Faculty Scientist II

Maine Medical Center Research Institute

DISCLOSURES

Source	Clinical Advisor	Employee	Research Funding
Maine Behavioral Healthcare		X	
NIMH, Simons Foundation, NLM Family Foundation			X
Camp Alsing	X		

AGENDA

- Aggression and self injury prevalence and impacts
- Multi-disciplinary approaches to serious challenging behaviors
- Studying the severely affected: The Autism Inpatient Collection (AIC)
- Novel approach to challenging behaviors
- Critical areas for further research

AGGRESSION AND SELF INJURY: A BIG PROBLEM

"My son is 19. Severely autistic. Can't speak. Getting more and more aggressive!!! Not only bites his wrist, he bangs walls and breaks things, and then comes after us. I can't keep locking myself in a room because he will bang the door down... No doctor in our area knows anything of what to do...Please, please I'm reaching out. There must be something for him...not fair for me and most certainly not fair to him! Cry for help!"

AGGRESSION PREVALENCE & IMPACTS

Up to 2/3 of youth with ASD develop aggression

 Kanne SM, Mazurek MO. J Autism Dev Disord. 2011

Aggression is one of the primary reasons youth with ASD use behavioral healthcare services

> Arnold LE, Vitiello B, McDougle C, et al. J Am Acad Child Adolesc Psychiatry. 2003

■ Families report that aggression is often of greater concern and negative impact than the core social and communicative deficits that define ASD

- Hodgetts S, et al. Focus Autism Other Dev Disabl. 2013 - Farmer CA, Aman MG. Res Autism Spectr Disord. 2011

Studies in ASD suggest that broadly-defined problem behaviors are heightened in ASD compared to typically developing (TD) and intellectually disabled (ID) samples; further, some ASD subgroups engage in persistent or increasing problem behaviors into adulthood.

PARTICULARLY FOR THE MINIMALLY VERBAL

- Inability to efficiently report distress makes aggression seem to occur "out of the blue."
 - Results in:
 - Caregivers unable to anticipate an aggressive outburst
 - Increasingly restrictive educational settings or exclusion
 - Inability to access community
 - High utilization of psychotropics, hospitalizations, out of home placements
 - Limits opportunities for real-time prevention (calming strategies, deescalation techniques, functional communication)
- The unmeasured effect of serious problem behaviors is a bending of the developmental trajectory <u>downward</u>

DEVELOPMENTAL DISORDERS SERVICE MAINE BEHAVIORAL HEALTHCARE

Spring Harbor Hospital DD Unit, Spring Harbor Academy, Day Treatment, Outpatient Clinic, Autism Research Team



Aggression is A Final Common Pathway Symptom

Aggression

Psychiatric Co-Morbidity

Behavioral Function & Reinforcement

Functional Communication Deficits

Side Effects

Dysregulated Sensory System Demands: abilities mismatch

Family Changes Medical Illness/Pain

Genetically Linked

Emotion Regulation

FOUNDATION OF TREATMENT

- Highly individualized behavioral plan with embedded communication and occupational therapy supports
- Targeted psychopharmacology
- Transfer of management skills to parents, local school, in-home staff

Multi-Disciplinary Treatment Team

- Child Psychiatry and Pediatrics
- Behavioral Psychologist
- Behavioral Coordinator (BCBA)
- Special Education
- Speech Pathologist
- Occupational Therapist
- Nursing
- Social Work
- Milieu Coordinator (OT)



THERAPEUTIC APPROACHES FOR CHALLENGING BEHAVIORS

- Applied Behavioral Analysis
- Psychotropic Medication

30% of functional behavioral assessment studies are inconclusive about behavioral function

- Derby KM, et al,1992, J Appl Behav A; Iwata BA, et al., J Appl Behav Anal. 1994; Vollmer TR, et al., J Appl Behav Anal. 1994


Significant side effects and inconsistent success

- Siegel M & Bealieu, A, JADD, 2011

- Communication strategies (AAC/Functional Communication)
- Treat Medical Problems
- Family treatment
- Parent Management Training (RUBI)
- Sensory regulation strategies
- Social skills / social cognitive strategies
- Psychotherapy approaches CBT / Emotion regulation

CHANGE IN BEHAVIORAL FUNCTIONING


Figure 1. Estimated Marginal Means of Abberant Behavior Checklist Irritability (ABC-I) Subscale Between Children with and without Autism Spectrum Disorder (ASD) Over Time



Siegel M, et al., JADD, July, 2014







AIC Autism Inpatient Collection

AIC RATIONALE

Knowledge & treatment options continue to lag for those with autism who are nonverbal, have an intellectual disability and/or display challenging behaviors; underrepresented in current, large data collections

- Communication: 30-50% do not develop functional verbal communication.
- Cognitive: 20-40% of individuals with Intellectual Disability (FSIQ<70)
- Behavioral: 20-30% with lifetime incidence of serious challenging behaviors

High volume of individuals - in a unique position to efficiently collect large amounts of data and improve understanding of this understudied portion of the ASD population

Inpatient setting an ideal platform to identify mechanisms underlying emotional and behavioral symptoms to inform treatment. Unique ability to study challenging behaviors *in situ* due to safety of inpatient environment and control over environmental factors

AUTISM INPATIENT COLLECTION




- Goals: Standardized assessment, description of the population, resource for all investigators, research platform for measure development, mechanistic studies and treatment studies.
 - To date, enrolled over 1000 probands and their biological parents
 - Rigorous core assessment battery and ASD diagnostic reliability
 - 48% minimally verbal, 42% Intellectual Disability (NVIQ <70; Molec Aut, 2015)
 - Whole exome sequencing to be performed 2018-19
 - Data available to approved investigators through SFARIBase
 - On-line community to facilitate recontacting through AUTISM INPATIENT



AIC ON SFARI BASE

SFARI Base



How to access SFARI Base and request SFARI resources: SFARI Base is a central database of phenotypic and genetic information about families affected by autism and other neurodevelopmental disorders, provided as part of the Simons Foundation Autism Research Initiative (SFARI). It contains data from the following cohorts:

- Simons Simplex Collection (SSC)
- Simons Variation in Individuals Project (Simons VIP)
- Simons Foundation Powering Autism Research for Knowledge (SPARK)
- Autism Inpatient Collection (AIC)




Predictors of Inpatient **Psychiatric Best practices for Outcomes** Hospitalization inpatient treatment **Problem** AIC Risk factors for **Psychotropic** behaviors and Methods **Self-injurious** Medication verbal ability behavior Use **Talking Expression of** about death Sleep and Trauma or suicide problem behavior Verbal ability and **Development of** psychiatric **Anxiety** the EDI symptoms **Parent stress** and problem Sleep and caregiver Health behavior stress disparities

AIC: CHANGE IN PROBLEM BEHAVIORS

ABC-I Scores, n=350


- Significant reduction in problem behavior scores from admission to discharge, and 2month follow-up (p<0.05)</p>
- Admission: 29.7(9.6)
- Discharge: 15.0(10.3)
- 2-Month Follow Up: 19.3(10.3)




- Pedersen K, et al., JADD, 2017

RYAN - THE NEED FOR NOVEL APPROACHES





IN HOSPITAL



RYAN - 23 YEAR OLD MAN



"Due to his inability to manage his emotions and communicate, he attacks us when he becomes agitated or stressed. This behavior is very unpredictable and often occurs with little to no warning signs. It has happened in our home, his school, in the community and sometimes while driving a car. It is so bad that we can no longer safely live in our home with Ryan."

-Wendi


PHYSIOLOGICAL AROUSAL AND CHALLENGING BEHAVIOR

In typically developing youth, greater ability to modulate physiological arousal is associated with fewer behavior problems. *Calkins SD. Dev Psychobilly. 1997; Porges, S.W. Dev Psyche; 1996

Association between physiological arousal and problem behavior in ASD

- -Cohen IL, et al., Assessing challenging behaviors in Autism Spectrum Disorders: Prevalence, rating scales, and autonomic indicators. In: *International handbook of autism and pervasive developmental disorders*. New York, NY: Springer; 2011;
- Groden J, et al, The impact of stress and anxiety on individuals with autism and developmental disabilities. In: *Behavioral Issues in Autism*. New York, NY: Springer; 1994;
- Romanczyk RG. Self-injurious behavior: Conceptualization, assessment, and treatment. In: *Advances in Learning & Behavioral Disabilities*. UK: Emerald Insight; 1986;
- Romanczyk RG, Lockshin S, O'Connor J. Psychophysiology and issues of anxiety and arousal. In: Self-injurious Behavior. New York, NY: Springer-Verlag; 1992


Hypothesis: Individual with ASD engages in a problem behavior <u>as an</u> <u>attempt</u> to alleviate distress and reestablish physiological homeostasis.




Time

Figure by Matthew Goodwin, PhD


CONCEPTUAL MODEL



TRANSFORMING THE TREATMENT OF CHALLENGING BEHAVIOR - REAL TIME PREDICTION




Pilot findings with 20 minimally-verbal inpatients



AUC values as a function of time to aggression, and signals used from past 3 minutes of data


- ~80 hrs of collection over ~70 naturalistic observation sessions
- Increasing accuracy as you increase
 # of signal streams utilized




ROC values for global and persondependent models using all combined features from the past 3 minutes to predict the next 1 minute

- Goodwin, MS, Ozdenizci, C,... Mazefsky, C, Siegel, M.,,, (in press). Predicting proximal aggression onset in minimally-verbal youth with autism spectrum disorder using preceding physiological signals. *Pervasive Health*.
- Ozdenizci, C... Mazefsky, C, Siegel, M,... & Goodwin, MS (in press). Time-series prediction
 of proximal aggression onset in minimally-verbal youth with autism spectrum disorder using
 physiological biosignals. Accepted to 40th International Conference of the IEEE Engineering
 in Medicine and Biology Society.

Staff Monitoring



Self Monitoring



CRITICAL UNMET RESEARCH NEEDS FOR THE SEVERELY AFFECTED

- Novel approaches to aggression and self injury that utilize objective, mechanistically-informed data
- Natural history of aggression and self-injury across the lifespan in ASD
- Validated, clinically practicable diagnostic tools for psychiatric co-morbidity
- Emotion regulation measurement and treatment
- Sleep biology and relationship to challenging behaviors
- Study of complex, real-world treatment packages (comparative effectiveness studies)
- Study of residential treatment and in-home behavioral services.

SUMMARY

- Individuals with ASD can develop serious behavioral challenges, which can become more impairing than the ASD itself.
- Parents report it is the <u>unpredictability</u> and <u>lack of warning</u> that causes the greatest impairment.
- Aggression and self injury are under-researched, and their underlying biology not well understood, particularly in those who are minimally verbal or have intellectual disability
- Pilot data from the AIC suggests that measurement of physiological arousal, combined with other data streams, can be used to predict the proximal onset of aggression.
- Critical need for novel research approaches to the assessment and treatment of aggression, self-injury and other problem behaviors, focusing on objective, biological mechanisms and measures.

ACKNOWLEDGEMENTS

Site Co-Investigators

- Carla Mazefsky, Ph.D, University of Pittsburgh
- Robin Gabriels, PsyD., Children's Colorado
- Giulia Righi, Ph.D. and Eric Morrow, MD, PhD, Brown University
- Craig Erickson, M.D. & Logan Wink, M.D., Cincinnatti Children's





Maine Medical Center Research Institute

- Susan Santangelo, Sc.D., Co-Investigator
- Kahsi A. Smith, Ph.D., Methodology/Statistics
- Mary Verdi, M.A., Project Manager
- Deanna Williams, B.A., Data Coordinator
- Christine Peura, B.A., Research Assistant
- Amy Stedman, B.A., Research Assistant
- Briana Taylor, PhD. Post-doctoral Fellow
- Donald St. Germain, M.D., Institute Director

Collaborators

- Matthew Goodwin, Ph.D., Northeastern University
- Matthew Lerner, Ph.D. & Kenneth Gadow, Ph.D & SUNY Stonybrook
- Louis Haggopian, Ph.D., Kennedy Krieger Institute
- Diane Williams, Ph.D., Duquesne University

THE Nancy Lurie Marks Family Foundation

Fostering Knowledge and Community for Autism and Beyond

ADDIRC Scientific Advisory Group

Connie Kasari, PhD, UCLA
Bryan King, MD, University of Washington
James McCracken, MD, UCLA
Christopher Mcdougle, MD, Harvard University
Lawrence Scahill, Ph.D, MSN, Emory University
Robert Schultz, Ph.D., University of Penn
Helen Tager-Flusberg, Ph.D., Boston University